

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
30 June 2005 (30.06.2005)

PCT

(10) International Publication Number
WO 2005/058370 A1

(51) International Patent Classification⁷: **A61K 49/00**

(21) International Application Number:
PCT/NO2004/000392

(22) International Filing Date:
17 December 2004 (17.12.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
20035681 18 December 2003 (18.12.2003) NO

(71) Applicant (for all designated States except US): **AMERSHAM HEALTH AS** [NO/NO]; Intellectual Property Dept., P.O. Box 4220 Nydalen, Nycoveien 1-2, N-0401 Oslo (NO).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KLAVENESS, Jo** [NO/NO]; Midtåsen 5, N-1166 Oslo (NO). **JOHANNESEN, Edvin** [NO/NO]; Amersham Health AS, P.O. Box 4220 Nydalen, Nycoveien 1-2, N-0401 Oslo (NO). **TOLLESHAUG, Helge** [NO/NO]; Amersham Health AS, P.O. Box 4220 Nydalen, Nycoveien 1-2, N-0401 Oslo (NO).

(74) Agents: **WULFF, Marianne, Weiby** et al.; Amersham Health AS, P.O. Box 4220 Nydalen, Nycoveien 1-2, N-0401 Oslo (NO).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: OPTICAL IMAGING CONTRAST AGENTS FOR IMAGING LUNG CANCER

(57) Abstract: The invention provides contrast agents for optical imaging of lung cancer in patients. The contrast agents may be used in diagnosis of lung cancer, for follow up of progress in disease development, for follow up of treatment of lung cancer and for surgical guidance. Further, the invention provides methods for optical imaging of lung cancer in patients.



WO 2005/058370 A1

OPTICAL IMAGING CONTRAST AGENTS FOR IMAGING LUNG CANCER

Field of the invention

The present invention provides contrast agents for optical imaging of lung cancer in
5 patients. The contrast agents may be used in diagnosis of lung cancer, for follow up
of progress in disease development, and for follow up of treatment of lung cancer.

The present invention also provides new methods of optical imaging of lung cancer in
patients, for diagnosis and for follow up of disease development and treatment of
10 lung cancer.

Description of related art

Lung cancer is the leading cause of cancer death worldwide. Approximately 25% of
all cancer deaths are attributed to lung cancer, and in USA alone, more than 160 000
15 new cases were diagnosed in year 2000 and more than 150 000 Americans died the
same year from lung cancer. Worldwide more than 1 million people died from lung
cancer in year 2000.

In general, the prognosis for patients with lung cancer is poor with a 5-year survival
20 rate of less than 15%. Nearly 90% of cases of lung cancer are attributed to cigarette
smoking.

Lung cancer can be divided into two distinct forms; small cell lung cancer (SCLC)
and non-small cell lung cancer (NSCLC). SCLC is without treatment the most
25 aggressive form of pulmonary tumours with median survival from diagnosis of two to
four months. Compared with other forms of lung cancer, SCLC is usually more
spread at time of diagnosis but is more responsive to chemotherapy and irradiation.
Chemotherapy of SCLC improves the survival time at least four to five fold. At the
time of diagnosis about one third of the patients have metastases in other organs.
30 Treatment of SCLC includes radiation therapy and chemotherapy. Typical drugs
used in treatment of SCLC include cisplatin, vincristine, doxorubicin, etoposide and
cyclophosphamide.

Non-small cell lung cancer (NSCLC) is a common terminology for various classes of
35 lung cancer including epidermoid carcinoma, adenocarcinoma and large cell
carcinoma. The disease can be treated in different ways depending on the stage of
disease at time of diagnosis. At an early stage the patient can undergo surgery as

this group of patients has the best prognosis. At a later stage the patients are usually treated with radiation therapy often in combination with chemotherapy. If the patients have metastases at the time of diagnosis they do not undergo surgery but are treated with radiation therapy or chemotherapy for palliation of symptoms from the primary
5 tumour.

Chemotherapeutic agents used for treatment of NSCLC include paclitaxel, docetaxel, topotecan, irinotecan, vinorelbine and gemcitabine.

- 10 Pulmonary function testing including spirometry and DLCO (diffusion capacity of the lung for carbon monoxide) is part of routine evaluation of lung cancer.

Conventional diagnostic staging of suspected lung malignancies involves chest radiography, bronchoscopy, CT of chest, ultrasound bone scans and MRI. MRI is
15 generally more sensitive than CT for diagnosis and staging of lung cancer. Recent advantages in diagnostic imaging of lung cancer include staging of the disease using PET and 18-fluorodeoxyglucose (FDG).

New bronchoscopic techniques like laser-induced fluorescence endoscope (LIFE)
20 bronchoscopy have the potential to improve the diagnosis of lung cancer.

Some methods have been described directed to measurements of lung function using light. US 4,646,750 (Williams) describes a method for detection of pulmonary inflammation using breath luminescence. US 5,227,308 (University of Hawaii) is
25 drawn to a method for assessing lung maturity using fluorescence from naphthalene-based probes. US 5,606,969 (Brigham & Women 's Hospital) relates to methods for measuring lung function using diffused light. US 4,534,360 (Williams) relates to a method for detection of lung cancer using breath luminescence.

30 The following documents describe compounds and methods for diagnosis for lung cancer. US 6,426,072 (Corixa) relates to compositions and methods for the therapy and diagnosis of lung cancer using lung tumour proteins and related substances. The document does not suggest imaging.

35 US 6,517,811 (Research Corporation Technologies) relates to compounds of cancer imaging and therapy including among others lung cancer. The compounds bind to a cell surface sigma receptor. Compounds including a radionuclide are described.

US 6,509,448 (Corixa) describes compositions and methods for the therapy and diagnosis of lung cancer. The compounds include polypeptides, polynucleotides encoding the polypeptides, antibodies, antigen presenting cells and immune system cells. The patent does not disclose optical imaging contrast agents.

US 6,509,316 (George Washington University) discuss compositions, methods and kits for treatment and diagnosis of lung cancer based on uteroglobin, for preventing/inhibiting metastasis of lung tumor cells. The patent does not describe optical imaging.

R. Baumgartner *et al*, J. Photochem Photobiol B 1996 36 169-74 studied the effect of inhaled 5-aminolevulinic acid to improve detection of early stage lung cancer. 5-aminolevulinic acid is not fluorescent, but is a biosynthetic precursor of the fluorescent protoporphyrin IX.

Lung cancer is still a challenge to diagnose and treat. There is a need for improved diagnostic methods, especially for diagnosis of lung cancer in an early stage with good reliability. Surprisingly, we have discovered that the use of optical imaging methods with new contrast agents fulfil these requirements.

Summary of the invention

The present invention provides an optical imaging contrast agent with affinity for an abnormally expressed biological target associated with lung cancer.

The invention is also described in the claims.

The following definitions will be used throughout the document:

Lung cancer tissue: Includes the two main forms small-cell lung cancer (SCLC) and non small-cell lung cancer (NSCLC), the latter including adenomas and squamous cell carcinomas. It further includes metastases to the lungs from other types of cancer.

Abnormally expressed target: A target that is either overexpressed or downregulated in lung cancer tissue.

Overexpressed target: A receptor, an enzyme or another molecule or chemical entity that is present in a higher amount in lung cancer tissue than in normal tissue.

5 Downregulated target: A receptor, an enzyme or another molecule or chemical entity that is present in a lower amount in lung cancer tissue than in normal tissue.

Detailed description of the invention

10 A first aspect of the present invention is an optical imaging contrast agent for imaging of lung cancer. By the term optical imaging contrast agent, or just contrast agent, we mean a molecular moiety used for enhancement of image contrast *in vivo* comprising at least one moiety that interacts with light in the ultraviolet, visible or near-infrared part of the electromagnetic spectrum.

15 The contrast agent has affinity for an abnormally expressed target associated with lung cancer.

Lung cancer tissue containing a downregulated target is identified by a low amount of bound imaging agent compared to normal tissue. In this situation, the amount of imaging agent should be less than 50 % of that in normal tissue, preferably less than
20 10 %.

Preferred contrast agents according to the invention, have affinity for an overexpressed target associated with lung cancer. Preferred targets are those targets that are more than 50 % more abundant in lung cancer tissue than in
25 surrounding tissue. More preferred targets are those targets that are more than two times more abundant in lung cancer tissue than in surrounding tissue. The most preferred targets are those targets that are more than 5 times more abundant in lung cancer tissue than in surrounding tissue.

30 In a further aspect of the invention, targets that are mutated in lung cancer tissue may be identified by lack of binding of an imaging agent that does bind to normal tissue; alternatively, the imaging agent might be directed specifically towards the mutated target, and binding to normal tissue would be minimal. The mutated target can be a protein in lung cancer tissue that is altered as a result of a germline or
35 somatic mutation, and including alterations resulting from differential splicing of RNA and changes in post-translational modifications, particularly glycosylation patterns, but not limited to these types of alterations.

Relevant groups of targets are receptors, enzymes, nucleic acids, proteins, lipids and other macromolecules as, for example, lipoproteins and glycoproteins. The targets may be located in the vascular system, in the extracellular space, associated with cell membranes or located intracellularly.

Preferred groups of targets are adhesion molecules and extracellular matrix proteins, antigens, proteins involved in cell cycle control and DNA repair, enzymes and inhibitors, hormones and hormone-related proteins, oncogens and receptors associated with lung cancer.

The following biological targets are overexpressed in lung cancer tissue and are preferred targets for contrast agents for optical imaging of lung cancer:

Adhesion molecules and extracellular matrix proteins

CD44, CD44v3, CD44v6, ED-B fibronectin, galectin-3, galectin-4, LGALS3 (Galectin) gene, P-selectin, liver-intestinal cadherin 17 and integrins, such as $\alpha_v\beta_3$ and $\alpha_v\beta_5$.

Antigens

CA 15.3, CA 72.4, cancer antigen 125 (CA125), CA19-9, carbohydrate antigen 549 (CA 549), carcinoembryonic antigen (CEA), CD105, CD24, CD34, chromogranin A antigens, ki-67, melanoma antigen E tumor-associated antigen, MUC1 (glycosylated mucin), oncoprotein 18, squamous cell carcinoma antigen (SCC), tissue polypeptide antigen (TPA), 5T4 oncofetal trophoblast glycoprotein, insulinoma-associated gene 1 product, FOS-related antigen 1, H/Le^y/Le^b.

Proteins involved in cell cycle control and DNA repair

K-ras, 34cdc2, Bax, bcl-2, Cdc 25A, cdc 25B, Cyclin B1, D1, E, cyclin D, p53, p27, pRb2/p130, retinoblastoma protein, telomerase, thyroid transcription factor 1, CDC6.

Enzymes and inhibitors

Cyclophilin A, alpha-1 protease inhibitor, arylamine N-acetyltransferase, Bcl2, carbonic anhydrase I and II, carbonic anhydrase-9, caspase-9 and -3, choline kinase, cyclo-oxygenase-2 (COX-2), CYP1A1, CYP2C40, cytidine deaminase, cytochrome P450, deoxycytidine deaminase, dual-specificity tyrosine-(Y)-phosphorylation regulated kinase 2 (DYRK 2), glutathione peroxidase, glutathione-S-transferase, GSTP1, GST-pi, helix-loop-helix ubiquitous kinase (CHUK), M2-PK (pyruvate

kinase), matrix metalloproteinases (MMPs), MMP-14, collagenase, MMP-9, Stromelysin-3 MutT homologue (hMTH1), an 8-oxo-dGTPase, myeloperoxidase, Neuron-specific enolase, phosphatidylinositol-3-kinase, prostaglandin E synthase, spermidine/spermine N1-acetyltransferase (SSAT), superoxide dismutase, thioether

5 S-methyltransferase, tyrosine kinase, urokinase plasminogen activator, ribonucleotide reductase, cystatin C, ERCC1 gene product, dopa decarboxylase, kallikrein 11, ornithine decarboxylase 1, cathepsin H, cathepsin L, farnesyl transferase, ribonucleotide reductase, tissue plasminogen activator, glutaminyl cyclase, pronapsin A, carbonyl reductase, leukotriene B4 12dehydrogenase,

10 thioredoxin reductase, glutathione peroxidase, glycylamide ribonucleotide formyltransferase (GARFT), thymidylate synthase, dihydrofolate reductase, carboxypeptidase E, proprotein convertase, protein kinase C-alpha, , ERCC1 gene product, ERCC2, hMLH1, hMSH2.

15 **Hormones and hormone-related proteins**

Arginine vasopressin, angiopoietin 1, angiopoietin 2, chromogranin A (CgA), CXC chemokines, ghrelin (growth hormone releasing peptide), interferon regulatory factor 1, macrophage migration inhibitory factor, pro-gastrin-releasing peptide (Pro-GRP), RANTES, vascular endothelial growth factor (VEGF), Insulin-like growth factor

20 binding protein 3 (IGFBP3), gastrin-releasing peptide, Cholecystokinin, neurotensin Insulin-like growth factor binding protein 3 (IGFBP3), calcitonin-related polypeptide and somatostatin.

Oncogenes

25 c-erbB-2, c-kit protein, EphA2 receptor tyrosine kinase, HER2/epidermal growth factor receptor (EGFR), HER-2/neu.

Receptors

Cholecystokinin A receptor, cholecystokinin B receptor, EGFR tyrosine kinase, epidermal growth factor receptor (EGFR), Notch3, TIE-2 precursor, SSR1 signal

30 sequence receptor- α , c-myc protein, Gastrin-releasing peptide receptor, neuromedin B receptor, bombesin receptor, neurotensin receptor, urokinase plasminogen activator receptor, vasopressin receptor, the angiopoietin receptors, vascular endothelial growth receptor (VEGFR), bradykinin receptor.

35

Other targets

Achaete scute homolog 1, alpha-1 PI2, alpha-adaptin, aryl hydrocarbon receptor, ataxia-telangectasia D-associated protein, AVP, BAG-1, beta-tubulin III, chromogranin-A, CYFRA, cytochrome b5, cytokeratin 19 fragment (Cyfra21-1), dickkopf homolog 1, differentiated embryo-chondrocyte, expressed gene 1 (DEC1) protein, dyskerin, eIF4E (translation initiation factor), epithelial mucin 1, ERK-1, ferritin, GRP, heat-shock proteins, hnRNP A2/B1, heterogeneous nuclear ribonucleoproteins, hnRNP B1, HSP70, HSP90, hypoxia-inducible factor (HIF) 1alpha, JAK-1, L523S (RNA-binding protein), MDR drug efflux/degradation, metallothionein, napsin A (TA02), NFAT1, p120, P16, proliferating cell nuclear antigen (PCNA), RAD21 homologue, retinoic acid receptor alpha, RhoA, ribonucleoprotein B1, S100 calcium-binding protein P, Solute carrier family 7, member 5, SpA, stanniocalcin 1, stathmin, surfactant proteins A, B, C and D, synaptophysin, thyroid transcription factor-1 (TTF-1), transmembrane protein 63 kD (ER/Golgi), UDG, uroplakin II, AKT, Ras, Ras-association domain family 1 (RASSF1A) protein, AFP, ALG-2, CC10, Kinin B1, MRP4, Nm23H1 gene.

Some targets are downregulated in lung cancer tissue and preferred targets are: Forkhead protein FREAC-1, Cadherin 5, Laminin β 1, Placenta copper monoamine oxidase, ABC3 ATP-binding cassette 3, Surfactant protein SP-C1, RAGE.

20

Among the more preferred targets for contrast agents for optical imaging of lung cancer are: Galectin-3, cancer antigen 125 (CA125), cathepsin L, MUC1, caspase-9 and -3, cyclo-oxygenase-2 (COX-2), glutathione-S-transferase (GST), the angiopoietin receptors, integrin α v β 3, vascular endothelial growth factor receptor (VEGF), HER2/epidermal growth factor receptor (EGFR), MDR, urokinase plasminogen activator receptor and cyclin D1.

25

The most preferred targets for contrast agents for optical imaging of lung cancer are cathepsin L, caspase-3, HER2/epidermal growth factor receptor (EGFR), urokinase plasminogen activator receptor and integrin α v β 3.

30

Generally, any targets that have been identified as possible targets for agents for treatment of lung cancer are potential targets also in optical imaging.

35 Small cell lung cancer (SCLC) synthesizes, and has receptors for several biologically active peptides that may be usable targets. The same receptors may not be relevant for non-small cell lung cancer (NSCLC).

The preferred contrast agents of the present invention are molecules with relatively low molecular weights. The molecular weight of preferred contrast agents is below 14 000 Daltons, preferably below 10000 Daltons and more preferably below 7000 Daltons.

The contrast agents, according to the present invention, are preferably comprised of a vector that has affinity for an abnormally expressed target in lung cancer tissue, and an optical reporter.

Thus viewed from one aspect the present invention provides a contrast agent of formula I:



wherein V is one or more vector moieties having affinity for one or more abnormally expressed target in lung cancer tissue, L is a linker moiety or a bond and R is one or more reporter moieties detectable in optical imaging.

The vector has the ability to direct the contrast agent to a region of lung cancer. The vector has affinity for the abnormally expressed target and preferably binds to the target. The reporter must be detectable in an optical imaging procedure and the linker must couple vector to reporter, at least until the reporter has been delivered to the region of lung cancer and preferably until the imaging procedure has been completed.

The vector can generally be any type of molecules that have affinity for the abnormally expressed target. The molecules should be physiologically acceptable and should preferably have an acceptable degree of stability. The vector is preferably selected from the following group of compounds: peptides, peptoids/peptidomimetics, oligonucleotides, oligosaccharides, lipid-related compounds like fatty acids, traditional organic drug-like small molecules, synthetic or semi-synthetic, and derivatives and mimetics thereof. When the target is an enzyme the vector may comprise an inhibitor of the enzyme or an enzyme substrate. The vector of the contrast agent preferably has a molecular weight of less than 10 000 Daltons, more preferably less than 4500 Daltons and most preferably less than 2500 Daltons, and hence does not include antibodies or internal image antibodies. In addition to

problems with immune reactions, long circulation time and limited distribution volume, many antibodies have an affinity for the receptor that is too low for use in imaging.

5 An optical imaging contrast agent comprising a vector having affinity for any of the preferred targets is a preferred embodiment of the invention.

Contrast agents having affinity for more than one abnormally expressed target related to the disease is an aspect of the invention. Such contrast agents can comprise two or more different vectors or molecular subunits that target two or more
10 different abnormally expressed targets.


Another possibility according to the present invention is that the contrast agent comprises one vector that is able to bind to more than one abnormally expressed target in lung cancer.
15

A contrast agent according to the present invention can also comprise more than one vector of same chemical composition that bind to the abnormally expressed biological target.

20 Some receptors are unique to endothelial cells and surrounding tissues. Examples of such receptors include growth factor receptors such as VEGF and adhesion receptors such as the integrin family of receptors. Peptides comprising the sequence arginine-glycine-aspartic acid (RGD) are known to bind to a range of integrin receptors. Such RGD-type peptides constitute one group of vectors for targets
25 associated with lung cancer.

Below are some examples of vectors having affinity for lung cancer-related abnormally expressed targets:

30 Vectors for cyclo-oxygenase-2 (COX-2): Arachidonic acid







Arachidonic acid [506-32-1] (Sigma A9673, A8798)

Arachidonic acid is the endogenous substrate for COX-2, and is an essential fatty acid and a precursor in the biosynthesis of prostaglandins..

Other vectors for COX-2 are exogenous compounds that bind to COX-2, for example
 5 so-called COX-2 inhibitors. The chemical classes of the main COX-2 inhibitors are shown in WO 02/07721.


Such vectors include:



10


Vectors for matrix metalloproteinases, such as for MMP-7:

Peptide sequence: Cys-Gly-Pro-Leu-Gly-Leu-Ala-Arg-OH




15


Vectors for mapping of tyrosine kinase activity of the epidermal growth factor receptor (EGFR):



5



10 Gefitinib (Iressa®):



These represent a group of kinase inhibitors and are analogues of ATP.

- 15 A wide variety of linkers can be used. The linker component of the contrast agent is at its simplest a bond between the vector and the reporter moieties. In this aspect the reporter part of the molecule is directly bound to the vector that binds to the abnormally expressed target. More generally, however, the linker will provide a mono- or multi-molecular skeleton covalently or non-covalently linking one or more
- 20 vectors to one or more reporters, e.g. a linear, cyclic, branched or reticulate molecular skeleton, or a molecular aggregate, with in-built or pendant groups which

bind covalently or non-covalently, e.g. coordinatively, with the vector and reporter moieties. The linker group can be relatively large in order to build into the contrast agent optimal size or optimal shape or simply to improve the binding characteristics for the contrast agent to the abnormally expressed target in lung cancer tissue.

5

Thus, linking of a reporter unit to a desired vector may be achieved by covalent or non-covalent means, usually involving interaction with one or more functional groups located on the reporter and/or vector. Examples of chemically reactive functional groups which may be employed for this purpose include amino, hydroxyl,

10

sulphydroxyl, carboxyl and carbonyl groups, as well as carbohydrate groups, vicinal diols, thioethers, 2-aminoalcohols, 2-aminothiols, guanidiny, imidazolyl and phenolic groups.

15

The reporter is any moiety capable of detection either directly or indirectly in an optical imaging procedure. The reporter can be a light scatterer (e.g. a coloured or uncoloured particle), a light absorber or a light emitter. More preferably the reporter is a dye such as a chromophore or a fluorescent compound. The dye part of the contrast agent can be any dye that interacts with light in the electromagnetic spectrum with wavelengths from the ultraviolet light to the near-infrared. Preferably,

20

the contrast agent of the invention has fluorescent properties.

25

Preferred organic dye reporters include groups having an extensive delocalized electron system, eg. cyanines, merocyanines, indocyanines, phthalocyanines, naphthalocyanines, triphenylmethines, porphyrins, pyrillium dyes, thiapyrillium dyes, squarylium dyes, croconium dyes, azulenium dyes, indoanilines, benzophenoxazinium dyes, benzothiaphenothiazinimum dyes, anthraquinones, naphthoquinones, indathrenes, phthaloylacridones, trisphenoquinones, azo dyes, intramolecular and intermolecular charge-transfer dyes and dye complexes, tropones, tetrazines, bis(dithiolene) complexes, bis(benzene-dithiolate) complexes, iodoaniline dyes, bis(S,O-dithiolene) complexes. Fluorescent proteins, such as green fluorescent protein (GFP) and modifications of GFP that have different absorption/emission properties are also useful. Complexes of certain rare earth metals (e.g., europium, samarium, terbium or dysprosium) are used in certain contexts, as are fluorescent nanocrystals (quantum dots).

35

Particular examples of chromophores which may be used include fluorescein, sulforhodamine 101 (Texas Red), rhodamine B, rhodamine 6G, rhodamine 19,

indocyanine green, Cy2, Cy3, Cy3B, Cy3.5, Cy5, Cy5.5, Cy7, Cy7.5, Marina Blue, Pacific Blue, Oregon Green 488, Oregon Green 514, tetramethylrhodamine, and Alexa Fluor 350, Alexa Fluor 430, Alexa Fluor 532, Alexa Fluor 546, Alexa Fluor 555, Alexa Fluor 568, Alexa Fluor 594, Alexa Fluor 633, Alexa Fluor 647, Alexa Fluor 660, Alexa Fluor 680, Alexa Fluor 700, and Alexa Fluor 750. The cyanine dyes are particularly preferred.

Particularly preferred are dyes which have absorption maxima in the visible or near-infrared region, between 400 nm and 3 μ m, particularly between 600 and 1300 nm.

The contrast agents according to the invention can comprise more than one dye molecular sub-unit. These dye sub-units can be similar or different from a chemical point of view. Preferred contrast agents have less than 6 dye molecular sub-units.

Several relevant targets for lung cancer are enzymes. A contrast agent for optical imaging of lung cancer for targeting an enzyme can be an enzyme contrast agent substrate that can be transformed to a contrast agent product possessing different pharmacokinetic and/or pharmacodynamic properties from the contrast agent substrate. This embodiment of the invention provides contrast agent substrates having affinity for an abnormally expressed enzyme, wherein the contrast agent substrate changes pharmacodynamic and/or pharmacokinetic properties upon a chemical modification into a contrast agent product in a specific enzymatic transformation, and thereby enabling detection of areas of disease upon a deviation in the enzyme activity from the normal. Typical differences in pharmacodynamic and/or pharmacokinetic properties can be binding properties to specific tissue, membrane penetration properties, protein binding and solubility properties.

Alternatively, if the abnormally expressed target for diagnosis of lung cancer is an enzyme, the contrast agent for optical imaging can be a dye molecule that directly binds to the enzyme. The contrast agent will have affinity for the abnormally expressed enzyme, and this may be used to identify tissue or cells with increased enzymatic activity.

In a further aspect of the invention, the contrast agent changes dye characteristics as a result of an enzymatic transformation. For example, a fluorescent dye reporter of the contrast agent is quenched (no fluorescence) by associated quencher groups,

until an enzymatic cleavage takes place, separating the dye from the quencher groups and resulting in fluorescence at the site of the abnormally expressed enzyme.

Another aspect of this part of the invention is that the dye may change colour, as e.g.
5 a change in absorption and/or emission spectrum, as a result of an enzymatic transformation.

If the abnormally expressed target for diagnosis of lung cancer is a receptor or another non-catalytic target, the contrast agent for optical imaging can bind directly to
10 the target and normally not change the dye characteristics.

The preferred contrast agents of the present invention are soluble in water. This means that the preferred contrast agents have a solubility in water at pH 7.4 of at least 1 mg/ml.
15

The contrast agents of the present invention can be identified by random screening, for example by testing of affinity for abnormally expressed targets of a library of dye labelled compounds either prepared and tested as single compounds or by preparation and testing of a mixture of compounds (a combinatorial approach).
20 Alternatively, random screening may be used to identify suitable vectors, before labelling with a reporter.

The contrast agents of the present invention can also be identified by use of technology within the field of intelligent drug design. One way to perform this is to use
25 computer-based techniques (molecular modelling or other forms of computer-aided drug design) or use of knowledge about natural and exogenous ligands (vectors) for the abnormally expressed targets. The sources for exogenous ligands can for example be the chemical structures of therapeutic molecules for targeting the same target. One typical approach here will be to bind the dye chemical sub-unit (reporter)
30 to the targeting vector so that the binding properties of the vector are not reduced. This can be performed by linking the dye at the far end away from the pharmacophore centre (the active targeting part of the molecule).

The contrast agents of the invention are preferably not endogenous substances
35 alone. Some endogenous substances, for instance estrogen, have certain fluorescent properties in themselves, but they are not likely to be sufficient for use in


optical imaging. Endogenous substances combined with an optical reporter however, fall within the contrast agents of the invention.

The contrast agent of the invention are intended for use in optical imaging. Any method that forms an image for diagnosis of disease, follow up of disease development or for follow up of disease treatment based on interaction with light in the electromagnetic spectrum from ultraviolet to near-infrared radiation falls within the term optical imaging. Optical imaging further includes all methods from direct visualization without use of any device and use of devices such as various scopes, catheters and optical imaging equipment, for example computer based hardware for tomographic presentations. The contrast agents will be useful with optical imaging modalities and measurement techniques including, but not limited to: luminescence imaging; endoscopy; fluorescence endoscopy; optical coherence tomography; transmittance imaging; time resolved transmittance imaging; confocal imaging; nonlinear microscopy; photoacoustic imaging; acousto-optical imaging; spectroscopy; reflectance spectroscopy; interferometry; coherence interferometry; diffuse optical tomography and fluorescence mediated diffuse optical tomography (continuous wave, time domain and frequency domain systems), and measurement of light scattering, absorption, polarisation, luminescence, fluorescence lifetime, quantum yield, and quenching.

Examples of contrast agents for optical imaging of lung cancer according to the invention, and potential synthesis of some of these, are shown below:


Contrast agents with affinity for mapping of COX-2:

1.



Wherein arachidonic acid, the endogenous substrate for COX-2, is linked to a reporter (R) via a linker (L).


2.



Wherein a COX-2 inhibitor derivative is linked to a reporter. R is any reporter according to the present invention; for example fluorescein, and L is a linker. For this example, giving a Rofecoxib-derivative, a possible synthesis is given.

Contrast agent for mapping of matrix metalloproteinase


The peptide vector (Cys-Gly-Pro-Leu-Gly-Leu-Ala-Arg) is linked to e.g. fluorescein (R) through a linker (L):



A synthesis is given in example 2.

Contrast agents for mapping of tyrosine kinase activity of the epidermal growth factor receptor (EGFR):

A suggested synthesis is given for preparation of a contrast agent comprising a vector with affinity for tyrosine kinase of the epidermal growth factor linked to a Cy5.5 reporter.



A further embodiment is the use of contrast agents of the invention for optical imaging of lung cancer, that is for diagnosis of lung cancer, for use in follow up the progress in lung cancer development, for follow up the treatment of lung cancer, or for surgical guidance.

In the context of this invention, diagnosis includes screening of selected populations, early detection, biopsy guidance, characterisation, staging and grading. Follow up of treatment includes therapy efficacy monitoring and long-term follow-up of relapse. Surgical guidance includes tumour margin identification during resection.

Still another embodiment of the invention is a method of optical imaging of lung cancer using the contrast agents as described.

Still another embodiment of the invention is a method of optical imaging for diagnosis, to follow up the progress of lung cancer development and to follow up the treatment of lung cancer, using a contrast agent as described.

5 One aspect of these methods is to administer the present contrast agents and follow the accumulation and elimination directly visually during surgery. Another aspect of these methods is to administer the present contrast agents and perform visual diagnosis through a bronchoscope.

10 Still another aspect of the present invention is to administer the present contrast agents and perform the image diagnosis using computerized equipment as for example a tomograph.

Still another embodiment of the invention is use of a contrast agent as described for
15 the manufacture of a diagnostic agent for use in a method of optical imaging of lung cancer involving administration of said diagnostic agent to an animate subject and generation of an image of at least part of said body, preferably the lungs or part of the lungs

20 Still another embodiment of the invention is pharmaceutical compositions comprising one or more contrast agents as described or pharmaceutically acceptable salts thereof for optical imaging for diagnosis of lung cancer, for follow up progress of lung cancer development or for follow up the treatment of lung cancer. The contrast agent of the present invention can be formulated in conventional pharmaceutical or
25 veterinary parenteral administration forms, e.g. suspensions, dispersions, etc., for example in an aqueous vehicle such as water for injections. The agent may also be formulated as an aerosol. Such compositions may further contain pharmaceutically acceptable diluents and excipients and formulation aids, for example stabilizers, antioxidants, osmolality adjusting agents, buffers, pH adjusting agents, etc. The most
30 preferred formulation is a sterile solution for intravascular administration or for direct injection into area of interest. Where the agent is formulated in a ready-to-use form for parenteral administration, the carrier medium is preferably isotonic or somewhat hypertonic.

35 The dosage of the contrast agents of the invention will depend upon the clinical indication, choice of contrast agent and method of administration. In general,

however dosages will be between 1 micro gram and 70 grams and more preferably between 10 micro grams and 5 grams for an adult human.

While the present invention is particularly suitable for methods involving parenteral
5 administration of the contrast agent, e.g. into the vasculature or directly into an organ
or muscle tissue, intravenous administration being especially preferred, it is also
applicable where administration is not via a parenteral route, e.g. where
administration is transdermal, nasal, sub-lingual or is into an externally voiding body
cavity, e.g. through the bronchi. The agent may be formulated as an aerosol for
10 administration by inhalation, or may be sprayed on directly during endoscopy. The
present invention is deemed to extend to cover such administration.

The following examples are illustrative only and not intended to be limiting. Other
features and advantages of the invention will be apparent from the detailed
15 description and from the claims.


Examples:**Example 1. Contrast agent for mapping of COX-2 activity. Synthesis of COX-2 ligand coupled to fluorescein.****Step 1**

2-Hydroxy-1-(4-methanesulfonylphenyl)ethanone is prepared from 2-bromo-1-(4-methanesulfonylphenyl)ethanone according to C. Puig *et al* in J.Med.Chem 2000, 43, 214-223.

Step 2

A solution of 2-hydroxy-1-(4-methanesulfonylphenyl) ethanone (1.50 g, 7 mmol) and fluorescein isocyanate isomer I (2.72 g, 7 mmol) is heated in DMF at 120°C for 5 hours.

The mixture is cooled, DMF evaporated off and acetic acid (25ml) is added. The mixture is refluxed for 10 hours. The acetic acid is evaporated and the resulting mixture is purified on silica using chloroform/methanol as eluent.

**Example 2. Contrast agent for mapping of matrix metalloproteinase (MMP). Synthesis of fluorescein-Cys-Gly-Pro-Leu-Gly-Leu-Leu-Ala-Arg-OH linker conjugate****Step 1**

The peptide component was synthesised on an ABI 433A automatic peptide synthesiser starting with Fmoc-Arg(Pmc)-wang resin on a 0.1 mmol scale using 1 mmol amino acid cartridges. The amino acids were pre-activated using HBTU before coupling. An aliquot of the peptide resin was then transferred to a clean round bottom flask an N-methyl morpholine (1 mmol) in DMF (5 ml) added followed by chloroacetyl chloride (1 mmol). The mixture was gently shaken until Kaiser test negative. The resin was extensively washed with DMF.

Step 2

5(6)-carboxyfluorescein (188 mg, 0.5 mmol) and dicyclohexylcarbodiimide (113 mg, 0.55 mmol) are dissolved in DMF (20 ml). The mixture is stirred for 2 hours and
5 cooled to 0°C. A solution of hexamethylenediamide (116 mg, 1 mmol) and DMAP (30 mg) in DMF is added and the mixture is stirred at ambient temperature for 72 hours. The solution is evaporated and the conjugate between carboxyfluorescein and hexamethylene-amine is isolated as monoamide by chromatography (silica, chloroform and methanol).


Step 3

The resin from step 1 is suspended in DMF (5 ml) and amide-amine conjugate from step 2 (0.5 mmol) pre-dissolved in DMF (5ml) containing triethylamine (0.5 mmol) is added. The mixture is heated to 50°C for 16 hours then excess reagents filtered off,
15 following extensive washing with DMF, DCM and diethyl ether then air drying. The product is treated with TFA containing TIS (5%), H₂O (5%), and phenol (2.5%) for 2 hours.

Excess TFA is removed *in vacuo* and the peptide is precipitated by the addition of diethyl ether. The crude peptide conjugate is purified by preparative HPLC C C-18, acetonitril, TFA, water).

Example 3. Contrast agent for binding to p53 oncoprotein**Step 1.** Synthesis of 2,2-bis(hydroxymethyl)-1-aza-bicyclo[2,2,2]octan-3-one.

25 3-quinuclidinone hydrochloride (Aldrich Q 190-5) (1 mmol) is dissolved in methanol-water (1:1, 30 ml). An aqueous solution of formaldehyde (37 %, 2.5 mmol) and sodium hydroxide (1.5 mmol) are added. The mixture is stirred for 12 hours at 50°C. The solvents are evaporated and the title compound isolated as free base using flash chromatography (silica, ethylacetate/chloroform, hexane).


Step 2.

5(6)-carboxyfluorescein (0.1 mmol) and dicyclohexyl carbodiimide (0.11 mmol) are dissolved in DMF. The mixture is stirred for 3 hours and cooled to 0 °C. A solution of 2,2-bis(hydroxymethyl)-1-azabicyclo[2,2,2] octane-3-one (0.5 mmol) and DMAP (10 mg) in DMF is added and the mixture is stirred at ambient temperature for 72 hours.

5 The solution is evaporated and the contrast agent is isolated by flash chromatography (silica, ethyl acetate/hexane).

Example 4 Contrast agent for mapping of tyrosine kinase activity of the epidermal growth factor.


10 **Step 1.** 4-[(3-bromophenyl)amino]-7-[N-(2-hydroxy-ethyl)-N-methylamino] pyrido [4,3-d] pyrimidine is prepared according to A.M. Thomson *et al* in J. Med. Chem. (1997) 40 3915-3925.




15 **Step 2.** 5(6)-carboxyfluorescein (1 mmol), dicyclohexylcarbodiimide (1.2 mmol) and DMAP (50 mg) are dissolved in DMF (30 ml). The mixture is stirred for 24 hours. A solution of the alcohol from step 1 (1 mmol) in DMF (5 ml) is added and the mixture is stirred for 3 days at ambient temperature. The fluorescein ester conjugate with the
20 alcohol vector is isolated by chromatography (silica, hexane/chloroform).

Example 5. Contrast agent for mapping of EGFR/erbB2 tyrosine kinase.


25 **Step 1.** N-[4-((3-bromophenyl)amino)quinazolin-7-yl]-acrylamide is prepared according to J. B. Smaill *et al* J. Med. Chem. (1999) 42 1803-1815.



Step 2. N-[4-((3-bromophenyl)amino)quinazolin-7-y]acrylamide from step 1 (1 mmol) and ethylenediamine (10 mmol) are dissolved in DMF (25 ml). The mixture is stirred at 50 °C for 12 hours. The solvent is evaporated off and the conjugate compound is isolated by flash chromatography (silica, hexane, chloroform, methanol).



Step 3. Cy7-NHS ester (0.5 mmol), the conjugate compound from step 2 (0.5 mmol) and N-methylmorpholine (70 mg) are dissolved in DMF (30 ml). The mixture is stirred at 40 °C for 3 days. The Cy7 amide conjugate is isolated by flash chromatography (silica, hexane, ethyl acetate, methanol).



Example 6. Inhalation formulation

The contrast agent from example 5 is filled into a powder inhalation device, e.g.

- 5 same type of device as the Pulmicort Turboinhaler ® from Astra Zeneca. The device contains 200 doses of 0.4 mg of the contrast agent. A contrast dose for diagnosis of lung cancer is typically 0.4 mg to 20 mg.

Example 7. Contrast agent with affinity for integrins: RGD peptide linked to Cy5.5


Step 1. Assembly of amino acids

The peptide sequence Asp-D-Phe-Lys-Arg-Gly was assembled on an Applied Biosystems 433A peptide synthesizer starting with 0.25 mmol Fmoc-Gly-SASRIN resin. An excess of 1 mmol pre-activated amino acids (using HBTU; O-Benzotriazol-1-yl-N,N,N',N'-tetramethyluronium hexafluorophosphate) was applied in the coupling steps. The cleavage of the fully protected peptide from the resins was carried out by treatment of the resin with three portions of 35 mL of 1 % trifluoroacetic acid (TFA) in

dichloromethane (DCM) for 5 minutes each. The filtrates containing the peptide was immediately neutralised with 2 % piperidine in DCM. The organics were extracted with water (3 x 100 mL), dried with MgSO_4 and evaporated *in vacuo*. Diethyl ether was added to the residue and the precipitate washed with ether and air-dried affording 30 mg of crude protected peptide. The product was analysed by analytical HPLC (conditions: Gradient, 20-70 % B over 10 min where A = $\text{H}_2\text{O}/0.1\%$ TFA and B = $\text{CH}_3\text{CN}/0.1\%$ TFA; flow, 2 mL/min; column, Phenomenex Luna 3μ 5 x 4.6 mm; detection, UV 214 nm; product retention time 7.58 min). Further product characterisation was carried out using electrospray mass spectrometry (MH^+ calculated, 1044.5; MH^+ found, 1044.4).

Step 2. N-C Cyclisation

c[-Asp-D-Phe-Lys-Arg-Gly-]




MW = 603.68
EM = 603.31
MF = C27H41N9O7

30 mg of the fully protected peptide, 16 mg of PyAOP, 4 mg of HOAt and 6 μL of N-methylmorpholine (NMM) were dissolved in dimethylformamide/DCM (1:1) and stirred over night. The mixture was evaporated *in vacuo* and diethyl ether added to the residue. The precipitate was washed with ether and air-dried. The crude cyclic fully protected peptide was treated with a solution of 25 mL TFA containing 5 % water, 5 % triisopropylsilane and 2.5 % phenol for two hours. TFA was evaporated *in vacuo* and diethyl ether added to the residue. The precipitate was washed with ether and air-dried. Purification by preparative RP-HPLC (0-30 % B over 40 min, where A = $\text{H}_2\text{O}/0.1\%$ TFA and B = $\text{CH}_3\text{CN}/0.1\%$ TFA, at a flow rate of 10 mL/min on a Phenomenex Luna 5μ C18 250 x 21.20 mm column) of the crude material afforded 2.3 mg pure product peptide. The pure product was analysed by analytical HPLC (conditions: Gradient, 0-15 % B over 10 min where A = $\text{H}_2\text{O}/0.1\%$ TFA and B = $\text{CH}_3\text{CN}/0.1\%$ TFA; flow, 2 mL/min; column, Phenomenex Luna 3μ 5 x 4.6 mm; detection, UV 214 nm; product retention time 6.97 min). Further product characterisation was carried out using electrospray mass spectrometry (MH^+ calculated, 604.3; MH^+ found, 604.4).

Step 3 . Conjugation of Cy5.5 to RGD peptide

c[-Asp-D-Phe-Lys(Cy5.5)-Arg-Gly-]




5

0.6 mg of the RGD peptide, 1.7 mg of Cy5.5 mono NHS ester and 5 μ L of NMM were dissolved in 1 mL of dimethylformamide (DMF) and the reaction mixture stirred for 2 hrs. Diethyl ether was added to the DMF solution and the blue precipitate washed with diethyl ether and air-dried affording 0.7 mg of crude RGD peptide conjugated to Cy5.5. The pure product was analysed by analytical HPLC (conditions: Gradient, 5-50 % B over 10 min where A = H₂O/0.1 % TFA and B = CH₃CN/0.1 % TFA; flow, 0.3 mL/min; column, Phenomenex Luna 3 μ 5 x 2 mm; detection, UV 214 nm; product retention time 8.32 min). Further product characterisation was carried out using electrospray mass spectrometry (MH⁺ calculated, 1502.5; MH⁺ found, 1502.6).

15

Example 8. Synthesis of 3-[(4'-Fluorobiphenyl-4-sulfonyl)-(1-hydroxycarbamoylcyclopentyl)amino]propionic acid (compound A) derivatised with Cy5.5 – contrast agent for binding to MMP

5



a) 1,11-Diazido-3,6,9-trioxaundecane

A solution of dry tetraethylene glycol (19.4 g, 0.100 mol) and methanesulphonyl chloride (25.2 g, 0.220 mol) in dry THF (100 ml) was kept under argon and cooled to 0 °C in an ice/water bath. To the flask was added a solution of triethylamine (22.6 g, 0.220 mol) in dry THF (25 ml) dropwise over 45 min. After 1 hr the cooling bath was removed and stirring was continued for 4 hrs. Water (60 ml) was added. To the mixture was added sodium hydrogencarbonate (6 g, to pH 8) and sodium azide (14.3 g, 0.220 mmol), in that order. THF was removed by distillation and the aqueous solution was refluxed for 24 h (two layers formed). The mixture was cooled and ether (100 ml) was added. The aqueous phase was saturated with sodium chloride. The phases were separated and the aqueous phase was extracted with ether (4 x 50 ml). Combined organic phases were washed with brine (2 x 50 ml) and dried (MgSO₄). Filtration and concentration gave 22.1 g (91%) of yellow oil. The product was used in the next step without further purification.

b) 11-Azido-3,6,9-trioxaundecanamine

To a mechanically, vigorously stirred suspension of 1,11-diazido-3,6,9-trioxaundecane (20.8 g, 0.085 mol) in 5% hydrochloric acid (200 ml) was added a solution of triphenylphosphine (19.9 g, 0.073 mol) in ether (150 ml) over 3 hrs at room temperature. The reaction mixture was stirred for additional 24 hrs. The phases were separated and the aqueous phase was extracted with dichloromethane (3 x 40 ml). The aqueous phase was cooled in an ice/water bath and pH was adjusted to ca

12 by addition of KOH. The product was extracted into dichloromethane (5 x 50 ml). Combined organic phases were dried (MgSO_4). Filtration and evaporation gave 14.0 g (88%) of yellow oil. Analysis by MALDI-TOF mass spectroscopy (matrix: α -cyano-4-hydroxycinnamic acid) gave a $\text{M}+\text{H}$ peak at 219 as expected. Further
5 characterisation using ^1H (500 MHz) and ^{13}C (125 MHz) NMR spectroscopy verified the structure.

c) Linking compound A to PEG(4)-N₃

To a solution of compound A (CP-471358, Pfizer, 41 mg, 87 μmol) in DMF (5 ml)
10 were added 11-azido-3,6,9-trioxaundecanamine (19 mg, 87 μmol), HATU (Applied Biosystems, 33 mg, 87 μmol) and DIEA (Fluka, 30 μl , 174 μmol). After one hour reaction time the mixture was concentrated and the residue was purified by preparative HPLC (column Phenomenex Luna C18(2) 5 μm 21.2 x 250 mm, solvents: A = water/0.1% TFA and B = acetonitrile/0.1% TFA; gradient 30-60% B over 60 min;
15 flow 10.0 ml/min, UV detection at 214 nm), giving 33.9 mg (59%) of product after lyophilisation. LC-MS analysis (column Phenomenex Luna C18(2) 3 μm 50 x 4.60 mm, solvents: A = water/0.1% TFA and B = acetonitrile/0.1% TFA; gradient 20-100% B over 10 min; flow 1 ml/min, UV detection at 214 nm, ESI-MS) gave a peak at 4.88 min with m/z 667.4 (MH^+) as expected.

d) Synthesis of compound A-PEG(4)-NH₂

To a solution of the PEG(4)-N₃ compound from c) (4.7 mg, 7.0 μmol) in methanol (4 ml) was added Pd/C (Koch-Light, ca 10 mg) added. The mixture was stirred at room temperature under hydrogen atmosphere (1 atm) for 10 min. The mixture was filtered
25 and concentrated. LC-MS analysis (column Phenomenex Luna C18(2) 3 μm 50 x 4.60 mm, solvents: A = water/0.1% TFA and B = acetonitrile/0.1% TFA; gradient 20-100% B over 10 min; flow 1 ml/min, UV detection at 214 nm, ESI-MS) gave a peak at 4.17 min with m/z 641.4 (MH^+) as expected. The product was used directly in the next step without further purification.

e) Conjugation of Cy 5.5

To a solution of the amine from d) (1.0 mg, 1.5 μmol) in DMF (0.2 ml) was added Cy 5.5-NHS (Amersham Biosciences, 1.0 mg, 1.0 μmol) and N-methylmorpholine (1 μl , 9 μmol). The reaction mixture was stirred for 48 h. MS analysis of the solution gave a
35 spectrum showing starting material and the conjugated product at m/z 1539.7 (M^+), expected 1539.4.

Example 9: Cy5-VEGF

Five micrograms of vascular endothelial growth factor (VEGF-121, cat.no. 298-VS/CH) (carrier-free, from R&D Systems) were dissolved in 19 μ l of 0.02 M borate buffer, pH 8.5. To this solution was added 2.5 nmol of the N-hydroxysuccinimide ester of a carboxylic acid derivative of Cy5 (Amersham Biosciences), dissolved in 5 μ l of the same buffer. The reaction mixture was incubated for one hour in the dark at room temperature. Unreacted dye was separated from the fluorescent protein derivative by centrifuging through a Micro-Spin 6 gel filtration column (Bio-Rad, exclusion limit about 6 kDa). The eluate fluoresced with excitation light at 646 nm, the emission being measured at 678 nm. The product was a fluorescent targeting molecule for the VEGF receptor.

Example 10: Cy5-TIMP-1

Five micrograms of tissue inhibitor of metalloproteinases-1 (TIMP-1, cat.no. 970-TM) (carrier-free, from R&D Systems) were dissolved in 25 μ l of 0.02 M borate buffer, pH 8.5. To this solution was added 2.5 nmol of the N-hydroxysuccinimide ester of a carboxylic acid derivative of Cy5 (Amersham Biosciences), dissolved in 5 μ l of the same buffer. The reaction mixture was incubated for one hour in the dark at room temperature. Unreacted dye was separated from the fluorescent protein derivative by centrifuging through a Micro-Spin 6 gel filtration column (Bio-Rad, exclusion limit about 6 kDa). The eluate fluoresced with excitation light at 646 nm, the emission being measured at 678 nm. The product was a fluorescent targeting molecule for matrix metalloproteinases.

Example 11: Fluorescein-TIMP-1

Five micrograms of tissue inhibitor of metalloproteinases-1 (TIMP-1, cat.no. 970-TM) (carrier-free, from R&D Systems) were dissolved in 25 μ l of 0.02 M borate buffer, pH 8.5. To this solution was added 2.5 nmol of the N-hydroxysuccinimide ester of a carboxylic acid derivative of fluorescein (Fluka), dissolved in 5 μ l of the same buffer. The reaction mixture was incubated for one hour in the dark at room temperature. Unreacted dye was separated from the fluorescent protein derivative by centrifuging through a Micro-Spin 6 gel filtration column (Bio-Rad, exclusion limit about 6 kDa). The eluate fluoresced with excitation light at 485 nm, the emission being measured at 538 nm. The product was a fluorescent targeting molecule for matrix metalloproteinases.

Example 12: Cy5-EGF

Sixty micrograms of epidermal growth factor (EGF, cat.no. 236-EG, 10 nmol) (from R&D Systems) were dissolved in 10 μ l of 0.02 M borate buffer, pH 8.5. To this solution was added 10 μ l buffer and 50 nmol of the N-hydroxysuccinimide ester of a carboxylic acid derivative of Cy5 (Amersham Biosciences). The reactive dye was dissolved in 5 μ l of the same buffer, mixed 1:1 with dioxan. The reaction mixture was incubated for one hour in the dark at room temperature. Unreacted dye was separated from the fluorescent protein derivative by centrifuging through a Micro-Spin 6 gel filtration column (Bio-Rad, exclusion limit about 6 kDa). The eluate, which was bright blue, fluoresced with excitation light at 646 nm, the emission being measured at 678 nm. The product was a fluorescent targeting molecule for the epidermal growth factor receptor.

Example 13: Cy7.5-EGF

Sixty micrograms of epidermal growth factor (EGF, cat.no. 236-EG, 10 nmol) (from R&D Systems) were dissolved in 10 μ l of 0.02 M borate buffer, pH 8.5. To this solution was added 10 μ l buffer and 50 nmol of the N-hydroxysuccinimide ester of a carboxylic acid derivative of Cy7.5 (Amersham Biosciences). The reactive dye was dissolved in 5 μ l of the same buffer, mixed 1:1 with dioxan. The reaction mixture was incubated for one hour in the dark at room temperature. Unreacted dye was separated from the fluorescent protein derivative by centrifuging through a Micro-Spin 6 gel filtration column (Bio-Rad, exclusion limit about 6 kDa). The eluate, which was dark green, fluoresced with excitation light at 700 nm, the emission being measured at 790 nm. The product was a fluorescent targeting molecule for the epidermal growth factor receptor.

Example 14: Fluorescein-EGF

Sixty micrograms of epidermal growth factor (EGF, cat.no. 236-EG, 10 nmol) (from R&D Systems) were dissolved in 10 μ l of 0.02 M borate buffer, pH 8.5. To this solution was added 10 μ l buffer and 50 nmol of the N-hydroxysuccinimide ester of a carboxylic acid derivative of fluorescein (Fluka), dissolved in 5 μ l of dioxan. The reaction mixture was incubated for one hour in the dark at room temperature. Unreacted dye was separated from the fluorescent protein derivative by centrifuging through a Micro-Spin 6 gel filtration column (Bio-Rad, exclusion limit about 6 kDa). The eluate, which was yellow, fluoresced with excitation light at 485 nm, the emission being measured at 538 nm. The product was a fluorescent targeting molecule for the epidermal growth factor receptor.

Claims:

1. An optical imaging contrast agent with affinity for an abnormally expressed biological target associated with lung cancer.

5

2. A contrast agent as claimed in claim 1 with molecular weight below 14 000 Daltons.

3. A contrast agent as claimed in claim 1 or 2 of formula I

10

V-L-R, (I)

wherein V is one or more vector moieties having affinity for an abnormally expressed target in lung cancer, L is a linker moiety or a bond and R is one or more reporter moieties detectable in optical imaging.

15

4. A contrast agent as claimed in any of claims 1 to 3 comprising a contrast agent substrate, wherein the target is an abnormally expressed enzyme, such that the contrast agent changes pharmacodynamic properties and/or pharmacokinetic properties upon a chemical modification from a contrast agent substrate to a contrast agent product upon a specific enzymatic transformation.

20

5. A contrast agent as claimed in any of claims 1 to 4 having affinity for any of the targets selected from galectin-3, cancer antigen 125 (CA125), cathepsin L, MUC1, caspase-9 and -3, cyclo-oxygenase-2 (COX-2), glutathione-S-transferase (GST), the angiopoietin receptors, integrin $\alpha v \beta 3$, vascular endothelial growth factor receptor (VEGF), HER2/epidermal growth factor receptor (EGFR), MDR, urokinase plasminogen activator receptor and cyclin D1.

25

6. A contrast agent as claimed in any of claims 3 to 5 wherein V is selected from peptides, peptoid moieties, oligonucleotides, oligosaccharides, fat-related compounds and traditional organic drug-like small molecules.

30

7. A contrast agent as claimed in any of claims 3-6 wherein R is a dye that interacts with light in the wavelength region from the ultraviolet to the near-infrared part of the electromagnetic spectrum.

35

8. A pharmaceutical composition for optical imaging of lung cancer comprising a contrast agent as defined in any of claims 1 to 7 together with at least one pharmaceutically acceptable carrier or excipient.

5 9. Use of a contrast agent as claimed in any of claims 1 to 7 for the manufacture of a diagnostic agent for use in a method of optical imaging of lung cancer involving administration of said diagnostic agent to an animate subject and generation of an image of at least part of said subject.

10 10. A method of optical imaging of lung cancer of an animate subject involving administering a contrast agent as defined in any of claims 1 to 7 to the subject and generating an optical image of at least a part of the subject to which said contrast agent has distributed.

15 11. Method as claimed in claim 10 for diagnosis of lung cancer, for follow up of the progress of lung cancer development, for follow up of treatment of lung cancer or for surgical guidance.

20 12. Use of a contrast agent as defined in any of claim 1 to 7 for optical imaging of lung cancer.

13. Use of a contrast agent as claimed in claim 12 for diagnosis of lung cancer, for follow up of the progress of lung cancer development, for follow up of treatment of lung cancer or for surgical guidance.

25

SEQUENCE LISTING

5 <110> Amersham Health AS

10 <120> Optical imaging contrast agents

<130> PN0398

15 <140> NO20035681

20 <141> 2003-12-18

<160> 2

25 <170> PatentIn version 3.1

30 <210> 1

<211> 5

35 <212> PRT

<213> Artificial Sequence

40 <220>

<223> Synthetic peptide

45 <400> 1

Asp Phe Lys Arg Gly
1 5

50 <210> 2

<211> 9

55 <212> PRT

<213> Artificial Sequence

60

<220>

<223> Synthetic peptide

5 <400> 2

Cys Gly Pro Leu Gly Leu Leu Ala Arg
1 5

10

INTERNATIONAL SEARCH REPORT

International Application No
PCT/NO2004/000392

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K49/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, EMBASE, CHEM ABS Data, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2001/055566 A1 (LUIKEN, M.D. GEORGE A ET AL) 27 December 2001 (2001-12-27) claims 1,6-11 paragraphs '0053! - '0055! paragraphs '0015! - '0017!, '0033! tables 1,2	1-3,6-13
X	WO 00/61194 A (INSTITUT FUER DIAGNOSTIKFORSCHUNG GMBH AN DER FREIEN UNIVERSITAET BERL) 19 October 2000 (2000-10-19) page 1, paragraph 1-3 page 3, paragraph 1-3 page 3, line 26 page 14, lines 13-30 claims 1,27 page 92, line 11 ----- -/--	1-3,6-13



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

9 May 2005

Date of mailing of the international search report

23/05/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Veronese, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/NO2004/000392

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 1 170 021 A (SHERING AKTIENGESELLSCHAFT) 9 January 2002 (2002-01-09) paragraphs '0026!', '0027! page 12, lines 6,9 paragraphs '0003! - '0006!; claim 1 -----	1-3,6-13
X	WO 01/91805 A (BRACCO RESEARCH USA; VON WRONSKI, MATHEW, A; MARINELLI, EDMUND, R; NUN) 6 December 2001 (2001-12-06) page 3, line 29 - page 4, line 28 sentences 10-21 See page 97, "BRU-317"; page 103, "BRU-326"; page 114, "BRU-346", page 120, "BRU-239". claims 1,10,47-53 -----	1-3,5-8
X	WO 98/47541 A (NYCOMED IMAGING AS; COCKBAIN, JULIAN, RODERICK, MICHAELSON; KLAVENESS,) 29 October 1998 (1998-10-29) See page 5, lines 1-4: vascular endothelial growth factors. page 70, paragraph 3 - page 86, paragraph 3 claims 1,4 -----	1-3,6-8
X	MARCHI-ARTZNER, VALERIE ET AL: "Adhesion of Arg-Gly-Asp (RGD) Peptide Vesicles onto an Integrin Surface: Visualization of the Segregation of RGD Ligands into the Adhesion Plaques by Fluorescence" LANGMUIR , 19(3), 835-841 CODEN: LANGD5; ISSN: 0743-7463, February 2003 (2003-02), XP002326372 the whole document -----	1-3,6,7
X	EP 0 800 831 A (DAIICHI PURE CHEMICALS CO. LTD; DAIICHI PURE CHEMICALS CO., LTD) 15 October 1997 (1997-10-15) page 2, line 50 page 7, line 37 example 12 page 17, lines 27-49 claim 1 -----	1,3
X	WO 00/71162 A (MALLINCKRODT INC) 30 November 2000 (2000-11-30) page 1, line 28 - page 2, line 2; claims; figures; examples ----- -/--	1-3,6-8

INTERNATIONAL SEARCH REPORT

International Application No
PCT/NO2004/000392

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>ACHILEFU S ET AL: "NOVEL RECEPTOR-TARGETED FLUORESCENT CONTRAST AGENTS FOR IN VIVO TUMOR IMAGING" INVESTIGATIVE RADIOLOGY, PHILADELPHIA, PA, US, vol. 35, no. 8, 2000, pages 479-485, XP000978923 the whole document</p>	1-13
Y	<p>WEINER RONALD E ET AL: "Radiolabeled peptides in the diagnosis and therapy of oncological diseases." APPLIED RADIATION AND ISOTOPES : INCLUDING DATA, INSTRUMENTATION AND METHODS FOR USE IN AGRICULTURE, INDUSTRY AND MEDICINE. NOV 2002, vol. 57, no. 5, November 2002 (2002-11), pages 749-763, XP002326772 ISSN: 0969-8043 paragraph '02.3!</p>	1-13
Y	<p>DATABASE MEDLINE 'Online! US NATIONAL LIBRARY OF MEDICINE (NLM), BETHESDA, MD, US; June 2003 (2003-06), TIAN FENG ET AL: "'Overexpression of COX-2 and its clinical significance in non-small cell lung cancer!'" XP002326773 Database accession no. NLM12895344 abstract & ZHONGHUA WAI KE ZA ZHI 'CHINESE JOURNAL OF SURGERY!. JUN 2003, vol. 41, no. 6, June 2003 (2003-06), pages 407-410, ISSN: 0529-5815</p>	1-13
Y	<p>DATABASE MEDLINE 'Online! US NATIONAL LIBRARY OF MEDICINE (NLM), BETHESDA, MD, US; 20 July 2001 (2001-07-20), KAJITA T ET AL: "The expression of vascular endothelial growth factor C and its receptors in non-small cell lung cancer." XP002326774 Database accession no. NLM11461086 abstract & BRITISH JOURNAL OF CANCER. 20 JUL 2001, vol. 85, no. 2, 20 July 2001 (2001-07-20), pages 255-260, ISSN: 0007-0920</p>	1-13

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/NO2004/000392

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	WO 2005/002293 A (VANDERBILT UNIVERSITY; MARNETT, LAWRENCE, J; TIMOFEEVSKI, SERGEI; PRUD) 6 January 2005 (2005-01-06) page 50, line 26 examples 4,19 claim 58 claim 1 See compounds 3, 18, 19, 24, 25. -----	1-4,6-13
E	WO 2005/030266 A (AMERSHAM HEALTH AS; KLAVENESS, JO; JOHANNESSEN, EDVIN; TOLLESHAUG, HELG) 7 April 2005 (2005-04-07) claims; examples -----	1-8
A	WO 01/89584 A (NYCOMED IMAGING AS; KLAVENESS, JO; TOLLESHAUG, HELGE; AMERSHAM HEALTH) 29 November 2001 (2001-11-29) page 21, paragraphs 2,3; claims; example 1 figure 2 -----	4
E	WO 2005/019247 A (BOARD OF REGENTS, THE UNIVERSITY OF TEXAS SYSTEM; LI, CHUN; KE, SHI; W) 3 March 2005 (2005-03-03) page 3, lines 19-23 page 5, lines 25-30 page 7, lines 29,30 figures 3-6,8 examples 2-4 claims 1,8-13,25,60 -----	1-3,6,7
E	WO 2005/003166 A (AMERSHAM HEALTH AS; CUTHBERTSON, ALAN; INDREVOLL, BAARD; SOLBAKKEN, MA) 13 January 2005 (2005-01-13) page 1, lines 1-10 page 2, line 25 page 5, lines 14-33 claims 1,12,13 -----	1-7

INTERNATIONAL SEARCH REPORT

International application No.
PCT/N02004/000392

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 10-13 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/NO2004/000392

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 2001055566	A1	27-12-2001	US 6284223 B1	04-09-2001
			US 6299860 B1	09-10-2001
			CA 2443927 A1	17-10-2002
			EP 1377221 A1	07-01-2004
			JP 2004527299 T	09-09-2004
			WO 02080778 A1	17-10-2002
			AU 764041 B2	07-08-2003
			AU 1308900 A	01-05-2000
			CA 2347461 A1	20-04-2000
			EP 1121158 A2	08-08-2001
			NZ 511175 A	26-09-2003
			WO 0021576 A2	20-04-2000
WO 0061194	A	19-10-2000	DE 19917713 A1	19-10-2000
			AT 259246 T	15-02-2004
			AU 769392 B2	22-01-2004
			AU 4291100 A	14-11-2000
			BG 105988 A	28-06-2002
			BR 0009658 A	15-01-2002
			CA 2368490 A1	19-10-2000
			CN 1351505 A	29-05-2002
			CZ 20013562 A3	15-05-2002
			DE 50005261 D1	18-03-2004
			DK 1176987 T3	14-06-2004
			EE 200100521 A	16-12-2002
			WO 0061194 A2	19-10-2000
			EP 1176987 A2	06-02-2002
			EP 1281405 A2	05-02-2003
			ES 2215641 T3	16-10-2004
			HR 20010833 A2	31-08-2004
			HU 0202990 A2	28-12-2002
			JP 2002541219 T	03-12-2002
			MX PA01010174 A	20-08-2003
			NO 20014911 A	06-12-2001
			NZ 514533 A	26-09-2003
			NZ 522135 A	28-05-2004
			NZ 522136 A	28-05-2004
			PL 351766 A1	16-06-2003
			PT 1176987 T	30-06-2004
			SK 14152001 A3	04-04-2002
			US 6630570 B1	07-10-2003
			ZA 200109238 A	10-02-2003
EP 1170021	A	09-01-2002	EP 1170021 A2	09-01-2002
WO 0191805	A	06-12-2001	AU 6669601 A	11-12-2001
			CA 2410887 A1	06-12-2001
			EP 1289565 A2	12-03-2003
			JP 2004500854 T	15-01-2004
			WO 0191805 A2	06-12-2001
			US 2002147136 A1	10-10-2002
WO 9847541	A	29-10-1998	AT 270116 T	15-07-2004
			AU 733477 B2	17-05-2001
			AU 4718297 A	22-05-1998
			AU 733495 B2	17-05-2001
			AU 4786697 A	22-05-1998
			AU 4786797 A	22-05-1998

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/N02004/000392

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 9847541	A		AU	4787097 A	22-05-1998
		AU	7068798 A	13-11-1998	
		BG	103438 A	31-01-2000	
		BG	103439 A	31-01-2000	
		BR	9712683 A	19-10-1999	
		BR	9713978 A	02-05-2000	
		CA	2269985 A1	07-05-1998	
		CA	2270120 A1	07-05-1998	
		CN	1440816 A	10-09-2003	
		CN	1238700 A	15-12-1999	
		CN	1234742 A	10-11-1999	
		CZ	9901494 A3	15-09-1999	
		DE	69824842 D1	05-08-2004	
		EP	1442751 A1	04-08-2004	
		EP	1007101 A2	14-06-2000	
		EP	0973552 A2	26-01-2000	
		EP	0991427 A2	12-04-2000	
		EP	0963209 A2	15-12-1999	
		EP	0977600 A2	09-02-2000	
		ES	2224379 T3	01-03-2005	
		WO	9818500 A2	07-05-1998	
		WO	9818501 A2	07-05-1998	
		WO	9818495 A2	07-05-1998	
		WO	9818498 A2	07-05-1998	
		WO	9847541 A1	29-10-1998	
		HU	0000357 A2	28-06-2000	
		JP	2001511765 T	14-08-2001	
		JP	2001503407 T	13-03-2001	
		JP	2002515889 T	28-05-2002	
		JP	2001502719 T	27-02-2001	
		JP	2002511845 T	16-04-2002	
		KR	2000052829 A	25-08-2000	
		KR	2000052830 A	25-08-2000	
		NO	991889 A	28-06-1999	
		NO	991890 A	28-06-1999	
		NZ	335596 A	27-10-2000	
		NZ	335799 A	24-11-2000	
		US	6610269 B1	26-08-2003	
		US	2004009122 A1	15-01-2004	
		US	2004141922 A1	22-07-2004	
		US	2005002865 A1	06-01-2005	
		US	2002102215 A1	01-08-2002	
EP 0800831	A	15-10-1997	AT	264694 T	15-05-2004
			AU	4497096 A	21-08-1996
			DE	69632244 D1	27-05-2004
			DE	69632244 T2	14-04-2005
			EP	0800831 A1	15-10-1997
			NO	973484 A	30-09-1997
			US	5968479 A	19-10-1999
			CA	2211470 A1	08-08-1996
			CN	1180316 A ,C	29-04-1998
			JP	9127115 A	16-05-1997
			WO	9623525 A1	08-08-1996
		WO 0071162	A	30-11-2000	US
	AU			4488800 A	12-12-2000
	CA			2373475 A1	30-11-2000

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/NO2004/000392

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0071162	A		EP 1178830 A2	13-02-2002
			JP 2003500367 T	07-01-2003
			WO 0071162 A2	30-11-2000
WO 2005002293	A	06-01-2005	US 2005002859 A1	06-01-2005
			WO 2005002293 A2	06-01-2005
WO 2005030266	A	07-04-2005	WO 2005030266 A2	07-04-2005
WO 0189584	A	29-11-2001	AU 7468301 A	03-12-2001
			EP 1283728 A2	19-02-2003
			JP 2003534297 T	18-11-2003
			WO 0189584 A2	29-11-2001
			US 2003170173 A1	11-09-2003
WO 2005019247	A	03-03-2005	US 2005069494 A1	31-03-2005
			WO 2005019247 A2	03-03-2005
WO 2005003166	A	13-01-2005	WO 2005003166 A1	13-01-2005